

Microscopic Modelling of the Optical Properties of Quantum-Well Semiconductor Lasers

*Stephan W. Koch
Department of Physics
Philipps University, Marburg/Germany*

- **OVERVIEW**

- Outline of Theory
- Gain/Absorption
- Luminescence
- Radiative and Auger Losses
- Nonequilibrium Effects

- **COLLABORATORS**

- A. Thränhardt, I. Kutzentsova, S. Becker, Marburg
- J. Hader, J. V. Moloney et al., Tucson
- W. Chow, Sandia

- **SUPPORT**

- AFOSR F49620-02-1-0380, DFG

Overview

simple models: easy to use

- often poorly controlled, ad hoc assumptions
- rely on major experimental input
- (semi-)empirical fits of experimental data
- many and adjustable parameters
- limited potential for predictions
-

microscopic theory: numerically demanding

- systematic approach based on fundamental physics
- controlled approximations
- predicts experiments
- no need for detailed experimental input
- wide range of applicability
- can be used as basis for controllably simplified models
-

Semiconductor Optics: Semiclassical Theory


MAXWELL'S WAVE EQUATION

$$\left[\frac{\partial^2}{\partial z^2} - \frac{n^2(z)}{c^2} \frac{\partial^2}{\partial t^2} \right] E = \mu_0 \frac{\partial^2}{\partial t^2} P$$

macroscopic optical polarization

semiconductors: Bloch basis

$$P = \sum_k d_{cv}^* P_k + c.c.$$




general theory: Haug/Koch, Quantum Theory of the Optical and Electronic Properties of Semiconductors, 4th ed., World Scientific (2004)

Semiconductor Density Matrix

two-band model, one QW subband

$$\begin{pmatrix} \langle a_k^\dagger a_k \rangle & \langle b_{-k} a_k \rangle \\ \langle a_k^\dagger b_{-k}^\dagger \rangle & \langle b_k^\dagger b_k \rangle \end{pmatrix} = \begin{pmatrix} f_k^c & P_k \\ P_k^* & f_k^v \end{pmatrix}$$



P_k ... e-h-pair transition amplitude


$f_k^{c,v}$... occupation probabilities

Hamiltonian


$$H = H_0 + H_{\text{Coul}} + H_{\text{dip}}$$

H_0 single particle (band structure)

H_{Coul} Coulomb interaction between carriers



H_{dip} dipole interaction with optical field



Calculation of Optical Response

Heisenberg equation of motion: $i\hbar \frac{\partial}{\partial t} \mathcal{O} = [\mathcal{O}, H]$

structure of resulting equations:

$$i\hbar \frac{\partial}{\partial t} \langle b_{-k} a_k \rangle = \varepsilon_k \langle b_{-k} a_k \rangle + \sum_{k', q \neq 0} V_q \langle a_{k'+q}^\dagger b_{q-k} a_{k'} a_k \rangle + \dots$$

infinite hierarchy of many-body correlations



systematic CLUSTER expansion


Schematics of Cluster Expansion

- two-point level (uncorrelated **single** particles)

$$\begin{aligned}
 i\hbar \frac{\partial}{\partial t} \langle \{1\}^\dagger \{1\} \rangle &= T [\langle \{1\}^\dagger \{1\} \rangle] + V [\langle \{2\}^\dagger \{2\} \rangle] \\
 &= T [\langle \{1\}^\dagger \{1\} \rangle] + V [\langle \{2\}^\dagger \{2\} \rangle_s] \\
 &\quad + V [\Delta \langle \{2\}^\dagger \{2\} \rangle]
 \end{aligned}$$

- four-point level (**two particle** correlations)

$$\begin{aligned}
 i\hbar \frac{\partial}{\partial t} \Delta \langle \{2\}^\dagger \{2\} \rangle &= T [\Delta \langle \{2\}^\dagger \{2\} \rangle] + V [\langle \{3\}^\dagger \{3\} \rangle] \\
 &= T [\Delta \langle \{2\}^\dagger \{2\} \rangle] + V [\langle \{3\}^\dagger \{3\} \rangle_s] \\
 &\quad + V [\langle \{3\}^\dagger \{3\} \rangle_D] \\
 &\quad + V [\Delta \langle \{3\}^\dagger \{3\} \rangle]
 \end{aligned}$$




Semiconductor Bloch Equations

all contributions beyond singlets

$$\left[i\hbar \frac{\partial}{\partial t} - \epsilon_k^e - \epsilon_k^h \right] P_k = [1 - f_k^e - f_k^h] \Omega_k + \frac{\partial}{\partial t} P_k|_{corr}$$

$$i\hbar \frac{\partial}{\partial t} f_k^a = -\Omega_k(t) P_k^* + \Omega_k^* P_k + \frac{\partial}{\partial t} f_k^a|_{corr}$$



HF field renormalization $\Omega_k(t) = d_{cv} E^{QW}(t) + \sum_{k'} V_{k-k'} P_{k'}(t)$

HF energy renormalization $\epsilon_k^a(t) = \epsilon_k^a - \sum_{k'} V_{k-k'} f_{k'}^a(t)$

- **nonlinearities**: phase-space filling, gap reduction, Coulomb enhancement
- **correlation contributions**: scattering, dephasing, screening


Second Born-Markov Approximation

- intraband Coulomb scattering
- phonon scattering


Born-Markov limit: Boltzmann equation

$$\frac{\partial}{\partial t} f_k^a(t)|_{corr} = \Sigma_k^{in,a}(t)[1 - f_k^a(t)] - \Sigma_k^{out,a}(t)f_k^a(t)$$

Coulomb scattering




phonon scattering



Coulomb Scattering I

$$\frac{\partial}{\partial t} f_k^a(t)|_{corr} = \Sigma_k^{in,a}(t)[1 - f_k^a(t)] - \Sigma_k^{out,a}(t)f_k^a(t)$$

scattering rates:




$$\frac{\partial}{\partial t} f_k^a|_{corr} \approx \sum_{k_1, k_2, k_3, b=e, h} |W|^2 f_{k_1}^a f_{k_2}^b [1 - f_{k_3}^b] [1 - f_k^a] - \dots$$

Coulomb Scattering II

$$\frac{\partial}{\partial t} f_k^a(t)|_{corr} = \Sigma_k^{in,a}(t)[1 - f_k^a(t)] - \Sigma_k^{out,a}(t)f_k^a(t)$$

quasi-equilibrium: $\frac{\partial}{\partial t} f_k^a|_{corr} = 0$
detailed balance

Fermi-Dirac distribution: $f_k^a \Rightarrow F_k^a = \frac{1}{e^{(E_k - \mu_a)/k_B T} + 1}$



Excitation Induced Dephasing

$$\left[i\hbar \frac{\partial}{\partial t} - \varepsilon_{\mathbf{k}}^e(t) - \varepsilon_{\mathbf{k}}^h(t) \right] P_{\mathbf{k}}(t) - [1 - f_{\mathbf{k}}^e(t) - f_{\mathbf{k}}^h(t)] \Omega_{\mathbf{k}}(t) \\ = i \left[\Gamma_{\mathbf{k}}(t) P_{\mathbf{k}}(t) + \sum_{k'} \Gamma_{\mathbf{k},\mathbf{k}'}(t) P_{\mathbf{k}'}(t) \right]$$


Re $\Gamma_{\mathbf{k}}(t)$... diagonal dephasing
⇒ generalized T_2 time
Im $\Gamma_{\mathbf{k}}(t)$... diagonal energy shift
⇒ generalized band-gap shift

Re $\Gamma_{\mathbf{k},\mathbf{k}'}(t)$... off-diagonal dephasing
Im $\Gamma_{\mathbf{k},\mathbf{k}'}(t)$... off-diagonal energy shift



- $\Gamma_{\mathbf{k}}(t)$ and $\Gamma_{\mathbf{k},\mathbf{k}'}(t)$ are calculated from all terms quadratic in the screened Coulomb interaction
- strong compensation between diagonal and off-diagonal terms

⇒ important for excitation saturation: Jahnke *et al.*, PRL 77, 5257 (1996).
and **gain calculations: Chow/Koch, Semiconductor-Laser Fundamentals, Springer Verlag (1999).**

Density Dependent Absorption




F. Jahnke, M. Kira, and S.W. Koch,
Z. Physik B **104**, 559 (1997)



- experiment: InGaAs/GaAs QW
- Khitrova, Gibbs, Jahnke, Kira, Koch, Rev. Mod. Phys. **71**, 1591 (1999)
- EID first observed in 4-wave mixing, Wang *et al.* PRL **71**, 1261 (1993)





Lineshape Problem





- $\Delta = (\hbar\omega - E_G)/E_B$
- gain of two-band bulk material
- nondiagonal scattering contributions $\rightarrow$ lineshape modification, no absorption below the gap

Hughes, Knorr, Koch, Binder, Indik, and Moloney
Solid State Comm. **100**, 555 (1996)

Optical Gain: Theory and Experiment




Wide Bandgap Materials: InGaN/GaN



- strong Coulomb effects: gapshift and broadening
- excitonic effects even at elevated densities and temperatures

Semiconductor Luminescence



$$H_{cf}^{qm} = \sum A(k, q) a_{k+q} b_{-k} B_q^\dagger + \text{h.c.}$$

where $A(k, q)$ is proportional to dipole matrix element and mode strength at the QW position

Semiconductor Luminescence Equations I

- photon operator dynamics

$$i\hbar \frac{\partial}{\partial t} B_{q_z q_{\parallel}} = \hbar \omega_q B_{q_z q_{\parallel}} + F_q \sum_k b_{-k} a_{k+q_{\parallel}} - F_q^* \sum_k a_{k-q_{\parallel}}^{\dagger} b_{-k}^{\dagger}$$

- polarization operator dynamics

$$i\hbar \frac{\partial}{\partial t} b_{-k} a_{k+q_{\parallel}} = \left(\tilde{E}_{k+q_{\parallel}}^e + \tilde{E}_k^h \right) b_{-k} a_{k+q_{\parallel}} \\ + \sum_{k' \neq k, k''} \left[V_{k'-k} b_{-k} \left(a_{k'+k''-k}^{\dagger} a_{k''} + b_{k'+k''-k}^{\dagger} b_{k''} \right) a_{k'+q_{\parallel}} + \dots \right] \\ - \sum_{q_{\parallel}'} a_{k+q_{\parallel}'}^{\dagger} \sum_{q_z} F_q^* B_{q_z q_{\parallel}'} a_{k+q_{\parallel}} + \dots$$

- Coulomb hierarchy problem (as in semiclassical theory)
- photon-carrier coupling hierarchy
- Cluster expansion to treat hierarchies on equal footing

Semiconductor Luminescence Equations II

$$i\hbar \frac{\partial}{\partial t} \Delta \langle B_q^\dagger B_{q'} \rangle = \hbar (\omega_{q'} - \omega_q) \Delta \langle B_q^\dagger B_{q'} \rangle + \sum_k (F_q \Pi_{k,q}^* - \text{c.c.})$$

photon assisted interband polarization $\Pi_{k,q} = \Delta \langle B_q^\dagger b_{-k} a_{k+q_{||}} \rangle$

$$\left(i\hbar \frac{\partial}{\partial t} - \tilde{E}_{k,q_{||}} + \hbar\omega_q \right) \Pi_{k,q} = \left(1 - f_{k+q_{||}}^e - f_k^h \right) \sum_{k'} V_{k-k'} \Pi_{k',q} \\ + \left. \frac{\partial \Pi_{k,q}}{\partial t} \right|_{col} + \Omega_{k,q}^{coh} + \Omega_{k,q}^{stim} + F_q S_{k,q}$$

incoherent source $S = \langle a^\dagger b^\dagger ab \rangle = f^e f^h + \Delta \langle a^\dagger b^\dagger ab \rangle$

coherent source $\Omega^{coh} \propto P, E$

feedback (cavity) Ω^{stim}

excitonic signatures $(1 - f^e - f^h) \sum V \Pi$

Quasi Stationary: Luminescence Elliott Equation

$$I_{\text{PL}}(\omega) \propto \text{Im} \left[\sum_{\nu} \frac{\phi_{\nu}^r(r=0)}{E_{\nu} - \hbar\omega - i\gamma_{\nu}} \sum_{k,k'} \left(\phi_{\nu}^l(k) \right)^{\star} \langle a_{k'}^{\dagger}, a_k b_k^{\dagger}, b_k \rangle \right]$$

- $\langle N_{\nu} \rangle_{\text{corr}} \propto \sum_{k,k'} \langle a_{k'}^{\dagger}, a_k b_k^{\dagger}, b_k \rangle = \langle N_{\nu} \rangle_{\text{plasma}} + \Delta \langle N_{\nu} \rangle$

with

- $\langle N_{\nu} \rangle_{\text{plasma}} \propto \sum_k f_k^e f_k^h$
- $\Delta \langle N_{\nu} \rangle$ „true“ excitons


- excitonic resonances independent of source term
- relative peak height depends on populations

Gain/Absorption and Luminescence

5nm $\text{Ga}_{0.8}\text{In}_{0.2}\text{As}/\text{GaAs}$ pin-MQW

gain: **theory**: 1.0, 1.125, 1.25, 1.375, 1.5, 1.625, 1.75, 1.875
experiment: 6.0, 6.5, 7.0, 7.5, 8.3, 9.0 [mA]

PL: **theory**: 0.17, 0.30, 0.40, 0.53, 0.68, 0.86 [$10^{12} / \text{cm}^2$]
experiment: 12, 16, 18, 21, 24 [mW]



Luminescence Analysis

Input:

- nominal structural parameters
- experimental luminescence spectra at two moderate intensities

Analysis:

calculate ideal, homogeneously broadened luminescence spectra
for given structure



compare to experimental luminescence



determine from deviations the inhomogeneous broadening, actual
structural parameters, quality of device



use optimized parameters to calculate gain/absorption spectra,
refractive index, differential gain, linewidth enhancement factor, ...


Ref.: Hader et al., IEEE Phot. Techn. Lett. **14**, 762 (2002)

Luminescence Analysis II

- sample with three quantum wells
- nominal composition: 5nm Ga(.8)In(.2)As/GaAs
- sandwiched between n and p doped buffer layers

experiment: 12, 16, 18, 21, 24 mW

theory: 0.1, 0.17, 0.23, 0.3, 0.4, 0.53, 0.68, 0.86 10^{12}cm^2



- needed: slight adjustment of composition
- inhomogeneous broadening

Luminescence Analysis

Input:

- nominal structural parameters
- experimental luminescence spectra at two moderate intensities

Analysis:

calculate ideal, homogeneously broadened luminescence spectra
for given structure



compare to experimental luminescence




determine from deviations the inhomogeneous broadening, actual
structural parameters, quality of device

Luminescence Analysis III

- sample with three quantum wells
- nominal composition: 5nm Ga(.8)In(.2)As/GaAs
- sandwiched between n and p doped buffer layers

experiment: 12, 16, 18, 21, 24 mW

theory: 0.1, 0.17, 0.23, 0.3, 0.4, 0.53, 0.68, 0.86 10^{12}cm^{-2}



- good fit to experiment yields actual composition
- less In (0.19) or thinner (- one monolayer)

Luminescence Analysis

Input:

- nominal structural parameters
- experimental luminescence spectra at two moderate intensities

Analysis:

calculate ideal, homogeneously broadened luminescence spectra
for given structure



compare to experimental luminescence



determine from deviations the inhomogeneous broadening, actual
structural parameters, quality of device




use optimized parameters to calculate gain/absorption spectra

Absorption/Gain

- using optimized material parameters
- experimental inhomogeneous broadening

theory: 0.5, 0.75, 1.0, 1.25, ..., 4.25, 4.5 [$10^{12}/\text{cm}^2$]




- other results:
- refractive index, alfa parameter, etc.

Test of Predicted Gain Spectra

- independently measured gain spectra
- NO additional fitting!

experiment: 5.0, 6.0, 6.5, 7.0, 7.5, 8.0, 8.6, 10.0 [mV]


theory: 1.0, 1.125, 1.25, 1.375, 1.5, 1.625, 1.75, 1.875 [10^{12}cm^2]



- very good theory/experiment agreement
- verification of microscopic theory
- predictive capabilities

Losses in Semiconductor Lasers

Classical Parametrization of Loss Current J_{loss} :




Problems with $\textcolor{green}{A}$, $\textcolor{red}{B}$, $\textcolor{blue}{C}$ - Parametrization:

- parameters only very roughly known and only for special cases;
depend on well- and barrier-materials, layer widths, temperatures, densities...
- **simple density-dependence far from reality**

Spontaneous Recombination Current: Examples


$$J_{spont} = eR_{spont} = e \int d\omega I_{PL}(\omega)$$



J_{spont} increases only linear with N at high densities

Auger Recombination I

$$J_{aug}^e = eR_{aug}^e = e \sum_{\mathbf{k}, \mathbf{i}} \frac{\partial}{\partial t} f_k^i|_{aug}$$



Inclusion of particle number non-conserving terms:


$$H_V = 2 \sum \left[a_{i_1, s, k+q}^\dagger b_{j_1, s', k'-q}^\dagger b_{j_2, -s, -k}^\dagger b_{j_3, s', k'} V_{s, s', s', -s}^{i_1, j_3, j_1, j_2}(q) \right. \\ \left. + b_{j_3, s, k+q}^\dagger b_{j_2, -s', -k'+q} b_{j_1, s, k} a_{i_1, s', k'} V_{-s', s, s, s'}^{j_2, j_1, j_3, i_1}(q) \right] - a \leftrightarrow b$$

**Impact
Ionization**

**Auger
Recombination**

Auger Recombination II

Quantum-Boltzmann scattering
in 2. Born-Markov approximation:




Impact Ionization

$$\frac{\partial}{\partial t} f_k^a|_{aug} = \frac{2\pi}{\hbar} \sum_{k_1, k_2, k_3, b=e,h} |W|^2 \left[f_{k_1}^b [1 - f_{k_2}^c] [1 - f_{k_3}^d] [1 - f_k^a] \right. \\ \left. - [1 - f_{k_1}^b] f_{k_2}^c f_{k_3}^d f_k^a \right] + \dots$$

Auger Recombination

Auger Recombination: Examples




J_{aug}

increases far less than cubic with N
sometimes even less than quadratic

Theory-Experiment Comparison I



Theoretical Procedure:

- calculate gain for various densities
- search for density that overcomes intrinsic losses (mirror losses) = threshold density
- calculate spontaneous emission and Auger recombination for this density
- **comparison with experimental data without adjustment**



(experimental data from
R.Fehse, et al., IEEE J. Sel.
Topics Quantum Electron.
8, 801 (2002))


Theory-Experiment Comparison II



experimental data:
A.F. Phillips et al.,
IEEE J. Sel. Topics
Quantum Electron. 5,
401 (1999))

Laser Modelling: VECSEL

Vertical External Cavity Surface Emitting Laser



Numerical simulation issues:

- optical pumping = input of carriers and kinetic energy
- nonequilibrium carrier distributions, nonequilibrium gain
- dynamic response

Nonequilibrium Gain Simulations

- Semiconductor Bloch Equations

$$\left[i\hbar \frac{\partial}{\partial t} - \epsilon_k^e - \epsilon_k^h \right] P_k = [1 - f_k^e - f_k^h] \Omega_k + \frac{\partial}{\partial t} P_k|_{corr}$$
$$i\hbar \frac{\partial}{\partial t} f_k^a = -\Omega_k(t) P_k^* + \Omega_k^* P_k + \frac{\partial}{\partial t} f_k^a|_{corr}$$

- Reduced Wave Equation

$$\frac{dE(t)}{dt} = -\gamma_c E(t) + \frac{i\omega\Gamma}{\epsilon_B V_{qw}} \sum_k \mu_k^* P_k(t)$$

Full numerical solution way too time consuming

APPROACH

- Calculate rates for scattering/relaxation/dephasing
- Solve coupled equations
- Interesting situations: pulsed and/or CW excitation

Relaxation Rate Model for Scattering


$$\frac{\partial}{\partial t} P_k|_{corr} = -\gamma_p P_k(t)$$

$$\frac{\partial}{\partial t} f_k^\alpha|_{corr} = -\gamma_{c-c} [f_k^\alpha - F_k^\alpha(\mu_p^\alpha, T_p)] - \gamma_{c-p} [f_k^\alpha - F_k^\alpha(\mu_l^\alpha, T_l)]$$

Parameters


- Relaxation rates are obtained by fits to microscopic calculations
- $F_k^\alpha(\mu_p^\alpha, T_p)$ is calculated considering energy and particle conservation
- $F_k^\alpha(\mu_l^\alpha, T_l)$ is calculated considering particle conservation
- Carrier-carrier scattering is an order of magnitude faster than carrier-phonon scattering
- Band structure is taken from microscopic calculations

Carrier Dynamics




10nm Ga_{0.764}In_{0.236}As QW between 20.3nm GaN_{0.007}As_{0.993},
Excitation by a 50fs pulse at 1.35eV at t=0

Effective Carrier-Carrier Scattering Times



- simple evaluation
- exponential fits work well



Electron-Hole Scattering Times



- Hole scattering faster because of larger effective mass
- Variation of scattering times with excitation energy (band structure vs availability of scattering partners)
- Increased scattering time for lower densities (e.g. for $1.5 \times 10^{12} \text{ cm}^{-2}$ $\tau_e = 354 \text{ fs}$, $\tau_h = 94 \text{ fs}$, instead of $\tau_e = 349 \text{ fs}$, $\tau_h = 87 \text{ fs}$ at $2.0 \times 10^{12} \text{ cm}^{-2}$)


Switch-On of Optically Pumped SCL

- dynamics of electron and hole distributions





- distinct kinetic hole in electron distribution
- small kinetic hole in hole distribution due to faster scattering time

Gain Dynamics





- initial absorption turns into gain
- transient gain overshoot

Carrier Temperature and Laser Intensity



Excitation Induced Temperature Change



- “hot” carriers (relative to the bandgap) can still cool down carrier distribution
- excitation energy = 0.61 eV $\Rightarrow$ Excitation at energy chemical potential $\Rightarrow \Delta T=0$
- hole chemical potential always in the bandgap

Summary

- predictive microscopic theory
- application to semiconductor laser gain media
- luminescence, Auger loss rates
- nonequilibrium gain, laser dynamics

MANY CHALLENGES:

- non-classical properties (quantum optics, quantum information science, ...)
- modified photonic environment (phot. x-tals, ...)
- microscopic modelling of disorder

References and more informations:

<http://www.physik.uni-marburg.de> (look for: semiconductor theory)

<http://www.nlcstr.com>